
FALL 2012 DRUGS!


HOW ARE WE DOING?


AISD Intervention Program


Bringing Parents, Community, and Schools together to build a safer and healthier future for our children


AISD Department of Guidance and Counseling


















Results of the 2012 District-wide Drug Use Survey


HOW DANGEROUS IT IS TO USE TOBACCO






Results of the 2012 District-wide Drug Use Survey

HOW DANGEROUS IT IS TO USE MARIJUANA





- More than 32% secondary students say that it is easy to procure alcohol, tobacco, and marijuana when they want it. This is down from 39% in 2010.
- Majority of parents strongly disapprove of their children using drugs
 - 80% Marijuana
 - 80% Tobacco
 - 67% Alcohol

- Alcohol use among 7th 12th grade students has decreased to 36% (40% in 2010); however, it is still high.
- 29% of seniors reported that alcohol was always served at parties they attended (32% in 2010); 29% also reported alcohol was never served at parties they attended.
- Approx. 8% (same in 2010) seniors reported attending school while drunk.
- 70% seniors report they would go to a friend for help if they had a drug problem. 59% report they would ask a parent for help.

- A larger percentage of 7th 12th grade girls (36.8%) reported using alcohol during the school year compared to 34.2% of boys.
- A greater percentage of 7th 12th grade boys reported using most other drugs compared to girls, with the exception of inhalants (boys 6.3%, girls 8.0%)
- Steroid use was reported by 1.1% of boys, and 1.1% of girls.
- Xanax (Alprazolam) was reportedly used by 3.2% (3.5% in 2010, 13.2% in 2008) of all secondary students, and Vicodin was used by 6.5% (3.1% in 2010). 2.4% report abusing an OTC decongestant. The highest reported prescription drug used was codeine, at 7.5%, followed closely by Vicodin at 6.5%.